

Study of boundary layer stability influence to WRF model forecasts quality for Siberia

Исследование влияния устойчивости пограничного слоя атмосферы на качество численных прогнозов модели WRF для территории Сибири


***Martynova Yu.V. (1), Zaripov R.B. (2),
Krupchatnikov V.N. (1,3)***

- (1) Siberian research hydrometeorological institute of Roshydromet, Novosibirsk, 630099, Sovetskaya, 30;
<http://sibnigmi.ru> (e-mail: FoxyJ13@gmail.com)
- (2) Hydrometeorological centre of Russia, Moskow, 123242, B. Predtechensky per., 11-13; <http://meteoinfo.ru>
- (3) Institute of computational mathematics and mathematical geophysics SB RAS, Novosibirsk, 630090, pr. Ak. Lavrent'eva, 6; <http://www.sccc.ru>

Hypothesis:

For conditions with a stable atmospheric boundary layer quality of the WRF ARW predictions is worse than for conditions with a unstable boundary layer.

Experiment



Periods:

20070206 – 20070216
 20070501 – 20070510
 20070705 – 20070714

20080110 – 20080131
 20080610 - 20080630

Model version:

WRF ARW 2.1

WRF ARW 3.0

Quality estimation:

RMS
 STDEV

Isobaric layer stability:

$$Ri = \frac{g}{\theta} \frac{\partial \theta / \partial z}{(\partial u / \partial z)^2 + (\partial v / \partial z)^2}$$

Conclusion

Results of this study corroborate the hypothesis about influence of atmospheric boundary layer state to WRF ARW predictions quality.

Thank you for your attention!

See you at my poster :-)